Personnal care after giving birth

Please indicate the topics for which you would like to receive additional information from our care team regarding your personal postnatal care. Please check the desired topics.

Postnatal bleeding and menstruation Postnatal bleeding gradually diminishes and	Pain relief You can take an analgesic (acetaminophen and/
changes colour. The bleeding lasts from two to six weeks. If you are not breastfeeding, your period will begin again in four to eight weeks after the birth. For breastfeeding	or ibuprofen) like the one you received in the hospital to relieve pain. These drugs are safe for breastfeeding mothers. Do not take them if you are allergic to them.
mothers, it may take several months for your period to return, even after you stop breastfeeding. Do not use tampons or vaginal douche. Change your sanitary pad regularly, at least every 4 hours.	Diet After your child is born, a varied, balanced diet is recommended according to Canada's Food Guide. Weightloss diets are not recommended during the postnatal
Uterine cramps Some mothers experience cramping after their child's birth. Cramping lasts about one week.	period or at any time while you are breastfeeding. Personal hygiene
Perineum (area between vagina and anus) The perineum area can remain sensitive for a while after the birth. If you have stitches, they should come out on	You may bathe or shower at any time after a vaginal birth. Bathing is allowed approximately 5 to 7 days after your C-section. Your tub should be clean with no bubble bath or oil.
their own within two weeks. Full healing of the tissue takes from four to six weeks.	Sleep and rest It is perfectly normal to feel tired after the birth
Urinary and bowel function You may experience some difficulty in urinating and having bowel movements for a few days after giving birth.	of a child. It may take several weeks before your energy level returns to normal. Regular rest periods are recommended; try to nap while your baby is sleeping.
If this situation persists, please discuss it with your doctor or midwife.	Postnatal exercise You can gradually resume exercising two weeks
Hemorrhoids Hemorrhoids often resolve on their own in less	after a vaginal birth and six weeks after a C-section.
than two weeks.	Sex and contraception Couples can begin having sex again whenever
Breast care For breastfeeding mothers, refer to My breastfeeding guide from Best start.	they decide the time is right. It is recommended that you discuss with your doctor or midwife the best method of contraception for you.
For mothers who are not breastfeeding, discuss caring for your breasts with your nurse, as your body will still produce milk in the days following delivery.	"Baby blues" and post-partum depression There is nothing unusual about feeling sad or overwhelmed by all of the feelings associated with your
C-section wound care and physical activity Your incision must be cleaned with mild, unscented	new role as a mother, or to have crying episodes. These feelings usually disappear without treatment in two
soap, rinsed and thoroughly dried. Remove the adhesive strips after 7 days if they have not peeled off on their own. Resume your physical activities gradually after a C-section.	weeks. The symptoms of post-partum depression are very similar to other types of depression, but thoughts centre more on the baby, i.e., you may focus too much or
Do not lift more than your baby during the first month. You can start driving your car again once you feel safe.	too little on your baby. Discuss your feelings with your doctor or midwife if they continue.

