

Postoperative instructions

Mastectomy

Please follow your surgeon's instructions if they differ from the following statements.

Your return home

Dressing:

- Remove your dressing for four days following the surgery.
- Remove the little white dressings (Steri-strips™) the eighth day following the surgery.
- You may shower. Your dressing is waterproof.
- No baths or pools are permitted for seven days.
- If there is a drain, please keep on a waterproof dressing until the removal of the drain.

Elastic (Tensor™):

- Keep the elastic on night and day for the first three days following surgery. This will help prevent pain and bruising. You may also adjust it if necessary.
- Remove the elastic when you shower. You should wear a sports bra which will help prevent any movement of the breast.
- If a tensor has not been placed on your chest, wear a sports bra from the first day of surgery on.

Physical activities

- It is important to move your arm and shoulder on the first day after your surgery (an exercise sheet will be given to you if you have had lymph nodes removed).
- No sports for two weeks.
- No driving for one week after your surgery or longer if you are taking pain medication.

Pain control and constipation prevention

- Take your prescribed medications regularly or take Tylenol™ and Advil™.
- Drink prune juice and/or take Metamucil™ if you feel constipated. If this is not effective, take one tablespoon of Milk of Magnesia daily or PEG 3350 (polyethylene glycol) such as Restoralax, Lax-a-day or Clearlax until you start having regular bowel movements.
- If this is not effective, take one tablespoon of milk of Magnesia daily until you start having regular bowel movements.
- If you have had a sentinel node biopsy, your urine will be green for approximately 24 to 36 hours and your skin will have a slightly greyish tint for approximately 12 hours.

Call your doctor or visit your nearest Emergency Department if you have:

- redness, swelling
- hardening or discharge (other than blood) from the incision
- fever and chills
- increase in pain
- pain and heat in your calf and/or
- sudden shortness of breath

Note: It is common for women's menstrual cycle to be irregular in the weeks following general anaesthesia.

To book your appointment, call your surgeon's office.

The office information was given your discharge instructions.

713 Montréal Rd. Ottawa ON

T 613.746.4621

hopitalmontfort.com